arXiv:1504.07364v1 [math.NT] 28 Apr 2015

Generators of the ring of weakly holomorphic modular functions for $\Gamma_1(N)$

Ja Kyung Koo and Dong Sung Yoon

Abstract

For a positive integer N divisible by $4, 5, 6, 7$ or 9, let $\mathcal{O}_{1,N}(\mathbb{Q})$ be the ring of weakly holomorphic modular functions for the congruence subgroup $\Gamma_1(N)$ with rational Fourier coefficients. We present explicit generators of the ring $\mathcal{O}_{1,N}(\mathbb{Q})$ over $\mathbb{Q}$ by making use of modular units which have infinite product expansions.

Contents

1 Introduction 1

2 Galois actions of modular function fields 3

3 Fricke functions and Siegel functions 4

4 Ring of weakly holomorphic modular functions for $\Gamma_1(N)$ 6

5 Generators of $\mathcal{O}_{1,N}(\mathbb{Q})$ 10

6 Fricke families of level N 13

1 Introduction

Let Γ be a congruence subgroup of $\mathrm{SL}_2(\mathbb{Z})$. A meromorphic modular function for Γ is called a *weakly holomorphic* if its poles (if any) are supported only at the cusps of Γ. For

---

2010 *Mathematics Subject Classification.* 11F03 (primary), 11F11, 11G16 (secondary).

Key words and phrases. modular functions, modular units

The second named author was supported by the National Institute for Mathematical Sciences, Republic of Korea.

a positive integer N, let

$$\begin{aligned}
\Gamma(N) &= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \mid \begin{bmatrix} a & b \\ c & d \end{bmatrix} \equiv \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \pmod{N} \right\}, \\
\Gamma_1(N) &= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \mid \begin{bmatrix} a & b \\ c & d \end{bmatrix} \equiv \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \pmod{N} \right\}, \\
\Gamma^1(N) &= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) \mid \begin{bmatrix} a & b \\ c & d \end{bmatrix} \equiv \begin{bmatrix} 1 & 0 \\ * & 1 \end{bmatrix} \pmod{N} \right\}.
\end{aligned}$$

By definition, we have $\Gamma \supset \Gamma(N)$ for some positive integer N. Furthermore, one can check that

$$\begin{bmatrix} N & 0 \\ 0 & 1 \end{bmatrix}^{-1} \Gamma(N) \begin{bmatrix} N & 0 \\ 0 & 1 \end{bmatrix} \supset \Gamma_1(N^2).$$

Hence if $f(\tau)$ is a modular function for $\Gamma(N)$, then $f(N\tau)$ is a modular function for $\Gamma_1(N^2)$. Here we note that the Fourier coefficients of $f(N\tau)$ coincide with those of $f(\tau)$. Therefore, the study of modular functions with respect to congruence subgroups is reduced to that of modular functions for $\Gamma_1(N)$.

For a rational vector $\mathbf{r} = \begin{bmatrix} r_1 \\ r_2 \end{bmatrix} \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$, we define the *Siegel function* $g_{\mathbf{r}}(\tau)$ on the complex upper half plane $\mathbb{H}$ by the following infinite product expansion

$$g_{\mathbf{r}}(\tau) = -q^{\frac{1}{2}\mathbf{B}_2(r_1)} e^{\pi i r_2 (r_1 - 1)} (1 - q^{r_1} e^{2\pi i r_2}) \prod_{n=1}^{\infty} (1 - q^{n+r_1} e^{2\pi i r_2})(1 - q^{n-r_1} e^{-2\pi i r_2}), \tag{1.1}$$

where $\mathbf{B}_2(X) = X^2 - X + 1/6$ is the second Bernoulli polynomial and $q = e^{2\pi i \tau}$. As is well known ([7] or [5, p.36]), it is a modular unit, that is, both zeros and poles are supported at the cusps of certain congruence subgroup.

Now, let $\mathcal{O}_{1,N}(\mathbb{Q})$ (resp. $\mathcal{O}_N^1(\mathbb{Q})$) be the ring of weakly holomorphic modular functions for $\Gamma_1(N)$ (resp. $\Gamma^1(N)$) with rational Fourier coefficients. When $N \equiv 0 \pmod{4}$, Eum et al ([2]) recently constructed explicit generators of $\mathcal{O}_N^1(\mathbb{Q})$ by means of Siegel functions and classify all Fricke families of such level N, which will be defined in §6.

In this paper we shall investigate how to generate the ring $\mathcal{O}_{1,N}(\mathbb{Q})$ for arbitrary N by improving their idea. We first construct generators of $\mathcal{O}_{1,N}(\mathbb{Q})$ over $\mathbb{Q}$ for $1 \leq N \leq 10$ and $N = 12$ by utilizing Siegel functions (Theorem 4.5). And, for given positive integers $m > 3$ and N divisible by m we will further present a primitive generator of $\mathcal{O}_{1,N}(\mathbb{Q})$ over $\mathcal{O}_{1,m}(\mathbb{Q})$, which is a Weierstrass unit (Theorem 5.2), from which we are able to determine generators of the ring $\mathcal{O}_{1,N}(\mathbb{Q})$ over $\mathbb{Q}$ when N is divisible by $4, 5, 6, 7$ or 9 (Corollary 5.3). Lastly, as byproduct, we can classify all Fricke families of such level N (Theorem 6.2).

Notation 1.1. The transpose of a matrix α is denoted by ${}^t\alpha$. If R is a ring with unity, $R^\times$ stands for the group of all invertible elements of R. For a positive integer N, let $\zeta_N = e^{2\pi i/N}$ be a primitive N-th root of unity.

2 Galois actions of modular function fields

In this section we shall describe the action of Galois groups between modular function fields.

For a positive integer N, let $\mathcal{F}_N$ be the field of meromorphic modular functions for $\Gamma(N)$ whose Fourier coefficients lie in the Nth cyclotomic field $\mathbb{Q}(\zeta_N)$.

Proposition 2.1. *$\mathcal{F}_N$ is a Galois extension of $\mathcal{F}_1$ and*

$$\mathrm{Gal}(\mathcal{F}_N/\mathcal{F}_1) \cong \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\} \cong G_N \cdot \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\},$$

where

$$G_N = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & d \end{bmatrix} \mid d \in (\mathbb{Z}/N\mathbb{Z})^\times \right\}.$$

More precisely, $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$ acts on $\mathcal{F}_N$ as follows:

(i) *The element $\begin{bmatrix} 1 & 0 \\ 0 & d \end{bmatrix} \in G_N$ acts on $h \in \mathcal{F}_N$ by*

$$h(\tau) = \sum_{n \gg -\infty} c_n q^{n/N} \longmapsto \sum_{n \gg -\infty} c_n^{\sigma_d} q^{n/N},$$

where $\sum_{n \gg -\infty} c_n q^{n/N}$ is the Fourier expansion of h and $\sigma_d \in \mathrm{Gal}(\mathbb{Q}(\zeta_N)/\mathbb{Q})$ satisfies $\zeta_N^{\sigma_d} = \zeta_N^d$.

(ii) *The element $\gamma \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$ acts on $h \in \mathcal{F}_N$ by*

$$h(\tau)^\gamma = (h \circ \widetilde{\gamma})(\tau),$$

where $\widetilde{\gamma}$ is a preimage of γ of the reduction map $\mathrm{SL}_2(\mathbb{Z}) \to \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$.

Proof. [6, Chapter 6, Theorem 3]. □

In a similar fashion as Proposition 2.1 (i), an element $\sigma \in \mathrm{Gal}(\mathbb{Q}_{\mathrm{ab}}/\mathbb{Q})$ induces an element of $\mathrm{Gal}(\mathcal{F}_N/\mathcal{F}_1)$ by applying σ to the Fourier coefficients. For $h \in \mathcal{F}_N$, we denote by h^σ the image of h under this automorphism.

3 Fricke functions and Siegel functions

For a lattice L in $\mathbb{C}$, we let

$$g_2(L) = 60 \sum_{\omega \in L \setminus \{0\}} \frac{1}{\omega^4}, \quad g_3(L) = 140 \sum_{\omega \in L \setminus \{0\}} \frac{1}{\omega^6}, \quad \Delta(L) = g_2(L)^3 - 27 g_3(L)^2.$$

And, we define the *Weierstrass $\wp$-function* (relative to L) and the *j-invariant* by

$$\begin{aligned} \wp(z; L) &= \frac{1}{z^2} + \sum_{\omega \in L \setminus \{0\}} \left(\frac{1}{(z-\omega)^2} - \frac{1}{\omega^2} \right) \quad (z \in \mathbb{C}), \\ j(\tau) &= 1728 \frac{g_2(\tau)}{\Delta(\tau)} \quad (\tau \in \mathbb{H}), \end{aligned}$$

where $g_2(\tau) = g_2([\tau, 1])$, $g_3(\tau) = g_3([\tau, 1])$ and $\Delta(\tau) = \Delta([\tau, 1])$. For a rational vector $\mathbf{r} = \begin{bmatrix} r_1 \\ r_2 \end{bmatrix} \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$, the *Fricke function* is defined by

$$f_{\mathbf{r}}(\tau) = -2^7 3^5 \frac{g_2(\tau) g_3(\tau)}{\Delta(\tau)} \wp(r_1 \tau + r_2; [\tau, 1]) \qquad (\tau \in \mathbb{H}).$$

Then $\mathbb{Q}[j(\tau)]$ is the ring of weakly holomorphic functions in $\mathcal{F}_1$ and we have

$$\mathcal{F}_N = \begin{cases} \mathbb{Q}(j(\tau)) & \text{if } N = 1 \\ \mathbb{Q}\left(j(\tau), f_{\mathbf{r}}(\tau) : \mathbf{r} \in \frac{1}{N}\mathbb{Z}^2 \setminus \mathbb{Z}^2\right) & \text{if } N > 1 \end{cases}$$

([6, Chapter 5, Theorem 2 and Chapter 6, §3]).

Proposition 3.1. *Let N (> 1) be an integer and $\mathbf{r} \in (1/N)\mathbb{Z}^2 \setminus \mathbb{Z}^2$.*

(i) *$f_{\mathbf{r}}(\tau)$ is weakly holomorphic and depends only on $\pm\mathbf{r} \pmod{\mathbb{Z}^2}$.*

(ii) *For $\alpha \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$ we obtain*

$$f_{\mathbf{r}}(\tau)^\alpha = f_{{}^t\alpha\mathbf{r}}(\tau).$$

Proof. [6, Chapter 6, §2-§3] □

The following two propositions describe the modularity criterion for Siegel functions and the relation between Fricke functions and Siegel functions.

Proposition 3.2. *Let N (> 1) be an integer and $\{m(\mathbf{r})\}_{\mathbf{r} = \left[\begin{smallmatrix} r_1 \\ r_2 \end{smallmatrix}\right] \in \frac{1}{N}\mathbb{Z}^2 \setminus \mathbb{Z}^2}$ be a family of integers such that $m(\mathbf{r}) = 0$ except for finitely many $\mathbf{r}$.*

(i) *If* $\sum_{\mathbf{r}} m(\mathbf{r}) \equiv 0 \pmod{12}$, *then for* $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ *we get*

$$\prod_{\mathbf{r}} g_{\mathbf{r}}(\tau)^{m(\mathbf{r})} \circ \gamma = \prod_{\mathbf{r}} g_{{}^t\gamma\mathbf{r}}(\tau)^{m(\mathbf{r})}.$$

(ii) *A finite product of Siegel functions*

$$\zeta \prod_{\mathbf{r}} g_{\mathbf{r}}(\tau)^{m(\mathbf{r})}$$

belongs to $\mathcal{F}_N$ *if*

$$\begin{aligned} &\sum_{\mathbf{r}} m(\mathbf{r})(Nr_1)^2 \equiv \sum_{\mathbf{r}} m(\mathbf{r})(Nr_2)^2 \equiv 0 \pmod{\gcd(2,N)\cdot N}, \\ &\sum_{\mathbf{r}} m(\mathbf{r})(Nr_1)(Nr_2) \equiv 0 \pmod{N}, \\ &\sum_{\mathbf{r}} m(\mathbf{r}) \cdot \gcd(12,N) \equiv 0 \pmod{12}. \end{aligned}$$

Here,

$$\zeta = \prod_{\mathbf{r}} e^{\pi i r_2(1-r_1)m(\mathbf{r})} \in \mathbb{Q}(\zeta_{2N^2}).$$

PROOF. See [4, Proposition 2.4], [5, Chapter 3, Theorem 5.2 and 5.3] (or [8]). □

PROPOSITION 3.3. *Let* $\mathbf{r}, \mathbf{s} \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$.

(i) $f_{\mathbf{r}}(\tau) = f_{\mathbf{s}}(\tau)$ *if and only if* $\mathbf{r} \equiv \pm\mathbf{s} \pmod{\mathbb{Z}^2}$.

(ii) *If* $\mathbf{r} \not\equiv \pm\mathbf{s} \pmod{\mathbb{Z}^2}$, *then we obtain*

$$f_{\mathbf{r}}(\tau) - f_{\mathbf{s}}(\tau) = 2^7 3^5 \frac{g_2(\tau)g_3(\tau)\eta(\tau)^4}{\Delta(\tau)} \cdot \frac{g_{\mathbf{r}+\mathbf{s}}(\tau)g_{\mathbf{r}-\mathbf{s}}(\tau)}{g_{\mathbf{r}}(\tau)^2 g_{\mathbf{s}}(\tau)^2},$$

where

$$\eta(\tau) = \sqrt{2\pi}\zeta_8 q^{1/24} \prod_{n=1}^{\infty}(1-q^n)$$

is the Dedekind η*-function which is a 24th root of* $\Delta(\tau)$.

PROOF. See [1, Lemma 10.4], [5, p.51].

□

REMARK 3.4. Let N (>1) be an integer and $\mathbf{r}, \mathbf{s}, \mathbf{r}', \mathbf{s}' \in (1/N)\mathbb{Z}^2 \setminus \mathbb{Z}^2$ such that $\mathbf{r} \not\equiv \pm\mathbf{s} \pmod{\mathbb{Z}^2}$ and $\mathbf{r}' \not\equiv \pm\mathbf{s}' \pmod{\mathbb{Z}^2}$. Then the function

$$\frac{f_{\mathbf{r}}(\tau) - f_{\mathbf{s}}(\tau)}{f_{\mathbf{r}'}(\tau) - f_{\mathbf{s}'}(\tau)} = \frac{g_{\mathbf{r}+\mathbf{s}}(\tau)g_{\mathbf{r}-\mathbf{s}}(\tau)}{g_{\mathbf{r}}(\tau)^2 g_{\mathbf{s}}(\tau)^2} \cdot \frac{g_{\mathbf{r}'}(\tau)^2 g_{\mathbf{s}'}(\tau)^2}{g_{\mathbf{r}'+\mathbf{s}'}(\tau)g_{\mathbf{r}'-\mathbf{s}'}(\tau)}$$

becomes a modular unit in $\mathcal{F}_N$ by Proposition 3.2 and 3.3 (ii), which is called a *Weierstrass unit* of level N.

4 Ring of weakly holomorphic modular functions for $\Gamma_1(N)$

For a positive integer N, let

$$\begin{aligned} X_1(N) &= \Gamma_1(N) \setminus \mathbb{H}^* \\ X^1(N) &= \Gamma^1(N) \setminus \mathbb{H}^*, \end{aligned}$$

where $\mathbb{H}^* = \mathbb{H} \cup \mathbb{Q} \cup \{i\infty\}$. Observe that $\Gamma^1(N)$ and $\Gamma_1(N)$ are conjugate in $\mathrm{SL}_2(\mathbb{R})$, that is,

$$\omega_N \Gamma^1(N) \omega_N^{-1} = \Gamma_1(N) \tag{4.1}$$

where $\omega_N = \begin{bmatrix} 1/\sqrt{N} & 0 \\ 0 & \sqrt{N} \end{bmatrix}$. Hence one can readily show that the map

$$\begin{array}{ccc} X_1(N) & \longrightarrow & X^1(N) \\ z & \longmapsto & \omega_N^{-1}(z) = Nz \end{array} \tag{4.2}$$

is a well-defined isomorphism between two modular curves. Now, let $\mathcal{F}_{1,N}(\mathbb{Q})$ (resp. $\mathcal{F}_N^1(\mathbb{Q})$) be the field of meromorphic modular functions for $\Gamma_1(N)$ (resp. $\Gamma^1(N)$) with rational Fourier coefficients. It then follows from (4.1) that the map

$$\begin{array}{ccc} \mathcal{F}_{1,N}(\mathbb{Q}) & \longrightarrow & \mathcal{F}_N^1(\mathbb{Q}) \\ h(\tau) & \longmapsto & (h \circ \omega_N)(\tau) = h(\tau/N) \end{array} \tag{4.3}$$

is an isomorphism. Furthermore the map (4.3) gives rise to a ring isomorphism

$$\mathcal{O}_{1,N}(\mathbb{Q}) \xrightarrow{\sim} \mathcal{O}_N^1(\mathbb{Q}). \tag{4.4}$$

PROPOSITION 4.1. *The genus of $X_1(N)$ is zero if and only if $1 \le N \le 10$ or $N = 12$. In this case, two sets of inequivalent cusps of $X_1(N)$ and $X^1(N)$ are as follows:*

N	inequivalent cusps of $X_1(N)$	inequivalent cusps of $X^1(N)$
1	$\{i\infty\}$	$\{i\infty\}$
2	$\{0, i\infty\}$	$\{0, i\infty\}$
3	$\{0, i\infty\}$	$\{0, i\infty\}$
4	$\{0, \frac{1}{2}, i\infty\}$	$\{0, 2, i\infty\}$
5	$\{0, \frac{1}{2}, \frac{2}{5}, i\infty\}$	$\{0, \frac{5}{2}, 2, i\infty\}$
6	$\{0, \frac{1}{2}, \frac{1}{3}, i\infty\}$	$\{0, 3, 2, i\infty\}$
7	$\{0, \frac{1}{2}, \frac{1}{3}, \frac{2}{7}, \frac{3}{7}, i\infty\}$	$\{0, \frac{7}{2}, \frac{7}{3}, 2, 3, i\infty\}$
8	$\{0, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{3}{8}, i\infty\}$	$\{0, 4, \frac{8}{3}, 2, 3, i\infty\}$
9	$\{0, \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{2}{9}, \frac{4}{9}, i\infty\}$	$\{0, \frac{9}{2}, 3, 6, \frac{9}{4}, 2, 4, i\infty\}$
10	$\{0, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{10}, i\infty\}$	$\{0, 5, \frac{10}{3}, \frac{5}{2}, 2, 4, 3, i\infty\}$
12	$\{0, \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{3}{4}, \frac{1}{5}, \frac{1}{6}, \frac{5}{12}, i\infty\}$	$\{0, 6, 4, 8, 3, 9, \frac{12}{5}, 2, 5, i\infty\}$

Proof. It is immediate from [3] and (4.2). □

From now on, we assume that $2 \leq N \leq 10$ or $N = 12$. Let $g_{1,N}(\tau)$ be the generator of the function field $\mathbb{C}(X_1(N))$ in the table ([4, Table 2]):

N	$g_{1,N}(\tau)$
2	$g_{\left[\begin{smallmatrix}1/2\\0\end{smallmatrix}\right]}(2\tau)^{12}$
3	$g_{\left[\begin{smallmatrix}1/3\\0\end{smallmatrix}\right]}(3\tau)^{12}$
4	$g_{\left[\begin{smallmatrix}2/4\\0\end{smallmatrix}\right]}(4\tau)^{8}g_{\left[\begin{smallmatrix}1/4\\0\end{smallmatrix}\right]}(4\tau)^{-8}$
5	$g_{\left[\begin{smallmatrix}2/5\\0\end{smallmatrix}\right]}(5\tau)^{5}g_{\left[\begin{smallmatrix}1/5\\0\end{smallmatrix}\right]}(5\tau)^{-5}$
6	$g_{\left[\begin{smallmatrix}3/6\\0\end{smallmatrix}\right]}(6\tau)^{3}g_{\left[\begin{smallmatrix}1/6\\0\end{smallmatrix}\right]}(6\tau)^{-3}$
7	$g_{\left[\begin{smallmatrix}2/7\\0\end{smallmatrix}\right]}(7\tau)^{2}g_{\left[\begin{smallmatrix}3/7\\0\end{smallmatrix}\right]}(7\tau)g_{\left[\begin{smallmatrix}1/7\\0\end{smallmatrix}\right]}(7\tau)^{-3}$
8	$g_{\left[\begin{smallmatrix}3/8\\0\end{smallmatrix}\right]}(8\tau)^{2}g_{\left[\begin{smallmatrix}1/8\\0\end{smallmatrix}\right]}(8\tau)^{-2}$
9	$g_{\left[\begin{smallmatrix}2/9\\0\end{smallmatrix}\right]}(9\tau)g_{\left[\begin{smallmatrix}4/9\\0\end{smallmatrix}\right]}(9\tau)g_{\left[\begin{smallmatrix}1/9\\0\end{smallmatrix}\right]}(9\tau)^{-2}$
10	$g_{\left[\begin{smallmatrix}3/10\\0\end{smallmatrix}\right]}(10\tau)g_{\left[\begin{smallmatrix}4/10\\0\end{smallmatrix}\right]}(10\tau)g_{\left[\begin{smallmatrix}1/10\\0\end{smallmatrix}\right]}(10\tau)^{-1}g_{\left[\begin{smallmatrix}2/10\\0\end{smallmatrix}\right]}(10\tau)^{-1}$
12	$g_{\left[\begin{smallmatrix}5/12\\0\end{smallmatrix}\right]}(12\tau)g_{\left[\begin{smallmatrix}1/12\\0\end{smallmatrix}\right]}(12\tau)^{-1}$

And, let $g_N^1(\tau) = g_{1,N}(\tau/N)$ be the function on the modular curve $X^1(N)$ induced from $g_{1,N}(\tau)$. Observe that $g_{1,N}(\tau)$ has rational Fourier coefficients, and so it belongs to $\mathcal{F}_{1,N}(\mathbb{Q})$ ([4, Theorem 6.7]).

Lemma 4.2. *Let N be as above. Then we have*

$$\begin{aligned} \mathcal{F}_{1,N}(\mathbb{Q}) &= \mathbb{Q}(g_{1,N}(\tau)), \\ \mathcal{F}_N^1(\mathbb{Q}) &= \mathbb{Q}(g_N^1(\tau)). \end{aligned}$$

Proof. If $\mathbb{C}(X_1(N)) = \mathbb{C}(S)$ for some subset $S \subset \mathcal{F}_{1,N}(\mathbb{Q})$, then $\mathcal{F}_{1,N}(\mathbb{Q}) = \mathbb{Q}(S)$ ([4, Lemma 4.1]). Therefore, the lemma follows from (4.3).

□

We shall first find generators of $\mathcal{O}_N^1(\mathbb{Q})$ in the above case. Since the genus of $X^1(N)$ is zero by Proposition 4.1 and (4.2), the map

$$\begin{array}{ccc} X^1(N) & \longrightarrow & \mathbb{P}^1(\mathbb{C}) \\ \tau & \longmapsto & [g_N^1(\tau) : 1] \end{array} \tag{4.5}$$

turns out to be an isomorphism between two compact Riemann surfaces ([4, Theorem 6.5]). Note that $g_N^1(i\infty) = \infty$ ([4, Table 3]). Let τ_0 be an inequivalent cusp of $X^1(N)$

other than $i\infty$. Then there exists $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ such that $\tau_0 = \gamma(i\infty)$ and we attain

$$g_N^1(\tau_0) = \lim_{\tau \to i\infty} (g_N^1 \circ \gamma)(\tau).$$

Since the function $g_N^1(\tau)$ satisfies the assumption of Proposition 3.2 (i), one can estimate the value $g_N^1(\tau_0)$ in a concrete way. For example, if $N = 5$ and $\tau_0 = 5/2$, then we derive

$$\begin{aligned} g_5^1(\tau_0) = \lim_{\tau \to i\infty} (g_5^1 \circ \left[\begin{smallmatrix} 5 & 2 \\ 2 & 1 \end{smallmatrix}\right])(\tau) &= \lim_{\tau \to i\infty} \frac{g_{\left[\begin{smallmatrix} 2 \\ 4/5 \end{smallmatrix}\right]}(\tau)^5}{g_{\left[\begin{smallmatrix} 1 \\ 2/5 \end{smallmatrix}\right]}(\tau)^5} \quad \text{by Proposition 3.2 (i)} \\ &= \lim_{\tau \to i\infty} \left(\frac{q^{13/12}\zeta_5^2(1 - q^{-1}\zeta_5^{-4})(1 - \zeta_5^{-4})}{q^{1/12}(1 - \zeta_5^{-2})} \right)^5 \quad \text{by (1.1)} \\ &= -2 - 10(\zeta_5 + \zeta_5^{-1}) - 5(\zeta_5^2 + \zeta_5^{-2}). \end{aligned}$$

Now, let

$$C_N = \{g_N^1(\tau_0) \mid \tau_0 \in \{\text{inequivalent cusps of } \Gamma^1(N) \text{ except } i\infty\}\}.$$

Then one can readily get the following table:

N	C_N
2	$\{0\}$
3	$\{0\}$
4	$\{16, 0\}$
5	$\{-2 - 5(\zeta_5 + \zeta_5^{-1}) - 10(\zeta_5^2 + \zeta_5^{-2}), -2 - 10(\zeta_5 + \zeta_5^{-1}) - 5(\zeta_5^2 + \zeta_5^{-2}), 0\}$
6	$\{8, -1, 0\}$
7	$\{4 + 3(\zeta_7 + \zeta_7^{-1}) + \zeta_7^2 + \zeta_7^{-2}, 4 + \zeta_7 + \zeta_7^{-1} + 3(\zeta_7^3 + \zeta_7^{-3}), 4 + 3(\zeta_7^2 + \zeta_7^{-2}) + \zeta_7^3 + \zeta_7^{-3}, 1, 0\}$
8	$\{3 + 2(\zeta_8 + \zeta_8^{-1}), -1, 3 - 2(\zeta_8 + \zeta_8^{-1}), 1, 0\}$
9	$\{2 + 2(\zeta_9 + \zeta_9^{-1}) + \zeta_9^2 + \zeta_9^{-2}, 2 + \zeta_9 + \zeta_9^{-1} + 2(\zeta_9^4 + \zeta_9^{-4}), -\zeta_3, -\zeta_3^2,$ $2 + 2(\zeta_9^2 + \zeta_9^{-2}) + \zeta_9^4 + \zeta_9^{-4}, 1, 0\}$
10	$\{2 + \zeta_{10} + \zeta_{10}^{-1} + \zeta_5 + \zeta_5^{-1}, \zeta_5^2 + \zeta_5^{-2}, 2 + \zeta_{10}^3 + \zeta_{10}^{-3} + \zeta_5^2 + \zeta_5^{-2}, \zeta_5 + \zeta_5^{-1}, 1, -1, 0\}$
12	$\{1 + \zeta_{12} + \zeta_{12}^{-1} + \zeta_6 + \zeta_6^{-1}, -1, -i, i, -\zeta_3, -\zeta_3^2, 1 - (\zeta_{12} + \zeta_{12}^{-1}) - (\zeta_3 + \zeta_3^{-1}), 1, 0\}$

The ordering in the set C_N corresponds to that of the set of inequivalent cusps of $X^1(N)$ in Proposition 4.1. Note that $C_N \subset \mathbb{Q}_{\mathrm{ab}}$ by (1.1), where $\mathbb{Q}_{\mathrm{ab}}$ is the maximal abelian extension of $\mathbb{Q}$.

Lemma 4.3. *Let $c \in C_N$ and $\sigma \in \mathrm{Gal}(\mathbb{Q}_{\mathrm{ab}}/\mathbb{Q})$. Then $c^\sigma \in C_N$.*

Proof. Since $\mathcal{F}_1 \subset \mathcal{F}_N^1(\mathbb{Q}) \subset \mathcal{F}_N$, it follow from Proposition 2.1 that $\mathcal{F}_N$ is a Galois extension of $\mathcal{F}_N^1(\mathbb{Q})$ whose Galois group is

$$\mathrm{Gal}(\mathcal{F}_N/\mathcal{F}_N^1(\mathbb{Q})) \cong G_N \cdot \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\} \mid \gamma \equiv \begin{bmatrix} 1 & 0 \\ * & 1 \end{bmatrix} (\mathrm{mod} N) \right\}. \tag{4.6}$$

Observe that $c = \lim_{\tau \to i\infty} (g_N^1 \circ \gamma)(\tau)$ for some $\gamma \in \mathrm{SL}_2(\mathbb{Z})$, and hence we achieve by Proposition 3.2 (i) and (1.1)

$$c^\sigma = \Big(\lim_{\tau \to i\infty} (g_N^1 \circ \gamma)(\tau)\Big)^\sigma = \lim_{\tau \to i\infty} (g_N^1 \circ \gamma)^\sigma(\tau).$$

Since $(g_N^1 \circ \gamma)^\sigma$ is a conjugate of g_N^1 over $\mathcal{F}_1$ and g_N^1 is G_N-invariant by (4.6), we see from Proposition 2.1 that

$$(g_N^1 \circ \gamma)^\sigma = g_N^1 \circ \gamma'$$

for some $\gamma' \in \mathrm{SL}_2(\mathbb{Z})$. Therefore, $c^\sigma \in C_N$. □

LEMMA 4.4. *Let $c \in \mathbb{C}$. Then $g_N^1(\tau) - c$ has neither zeros nor poles on $\mathbb{H}$ if and only if $c \in C_N$.*

PROOF. It is immediate from the isomorphism (4.5). □

For each $c \in C_N$, we let $f_{N,c}(x) \in \mathbb{Q}[x]$ be the minimal polynomial of c over $\mathbb{Q}$.

THEOREM 4.5. *Let N be an integer such that $2 \leq N \leq 10$ or $N = 12$. Then we have*

$$\begin{aligned} \mathcal{O}_{1,N}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,N}(\tau), f_{N,c}(g_{1,N}(\tau))^{-1}\big]_{c \in C_N} \\ \mathcal{O}_N^1(\mathbb{Q}) &= \mathbb{Q}\big[g_N^1(\tau), f_{N,c}(g_N^1(\tau))^{-1}\big]_{c \in C_N}. \end{aligned}$$

More precisely, we attain

$$\begin{aligned}
\mathcal{O}_{1,2}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,2}(\tau), g_{1,2}(\tau)^{-1}\big] \\
\mathcal{O}_{1,3}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,3}(\tau), g_{1,3}(\tau)^{-1}\big] \\
\mathcal{O}_{1,4}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,4}(\tau), g_{1,4}(\tau)^{-1}, (g_{1,4}(\tau) - 16)^{-1}\big] \\
\mathcal{O}_{1,5}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,5}(\tau), g_{1,5}(\tau)^{-1}, (g_{1,5}(\tau)^2 - 11g_{1,5}(\tau) - 1)^{-1}\big] \\
\mathcal{O}_{1,6}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,6}(\tau), g_{1,6}(\tau)^{-1}, (g_{1,6}(\tau) - 8)^{-1}, (g_{1,6}(\tau) + 1)^{-1}\big] \\
\mathcal{O}_{1,7}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,7}(\tau), g_{1,7}(\tau)^{-1}, (g_{1,7}(\tau) - 1)^{-1}, (g_{1,7}(\tau)^3 - 8g_{1,7}(\tau)^2 + 5g_{1,7}(\tau) + 1)^{-1}\big] \\
\mathcal{O}_{1,8}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,8}(\tau), g_{1,8}(\tau)^{-1}, (g_{1,8}(\tau) + 1)^{-1}, (g_{1,8}(\tau) - 1)^{-1}, (g_{1,8}(\tau)^2 - 6g_{1,8}(\tau) + 1)^{-1}\big] \\
\mathcal{O}_{1,9}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,9}(\tau), g_{1,9}(\tau)^{-1}, (g_{1,9}(\tau) - 1)^{-1}, (g_{1,9}(\tau)^2 - g_{1,9}(\tau) + 1)^{-1}, \\
&\qquad (g_{1,9}(\tau)^3 - 6g_{1,9}(\tau)^2 + 3g_{1,9}(\tau) + 1)^{-1}\big] \\
\mathcal{O}_{1,10}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,10}(\tau), g_{1,10}(\tau)^{-1}, (g_{1,10}(\tau) + 1)^{-1}, (g_{1,10}(\tau) - 1)^{-1}, (g_{1,10}(\tau)^2 + g_{1,10}(\tau) - 1)^{-1}, \\
&\qquad (g_{1,10}(\tau)^2 - 4g_{1,10}(\tau) - 1)^{-1}\big] \\
\mathcal{O}_{1,12}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,12}(\tau), g_{1,12}(\tau)^{-1}, (g_{1,12}(\tau) + 1)^{-1}, (g_{1,12}(\tau) - 1)^{-1}, (g_{1,12}(\tau)^2 + 1)^{-1}, \\
&\qquad (g_{1,12}(\tau)^2 - g_{1,12}(\tau) + 1)^{-1}, (g_{1,12}(\tau)^2 - 4g_{1,12}(\tau) + 1)^{-1}\big].
\end{aligned}$$

Proof. Since every zero of $f_{N,c}(x)$ lies in the set C_N by Lemma 4.3, $f_{N,c}(g_N^1(\tau))$ is a modular unit in $\mathcal{F}_N^1(\mathbb{Q})$ by Lemma 4.4. This shows that $\mathcal{O}_N^1(\mathbb{Q}) \supseteq \mathbb{Q}\big[g_N^1(\tau), f_{N,c}(g_N^1(\tau))^{-1}\big]_{c\in C_N}$.

Conversely, let $h(\tau) \in \mathcal{O}_N^1(\mathbb{Q})$. By Lemma 4.2 we can write

$$h(\tau) = \frac{P(g_N^1(\tau))}{Q(g_N^1(\tau))}$$

for some polynomials $P(x), Q(x) \in \mathbb{Q}[x]$ which are relatively prime. Suppose that $Q(x)$ has a zero $c_0 \in \overline{\mathbb{Q}} \setminus C_N$, where $\overline{\mathbb{Q}}$ is the algebraic closure of $\mathbb{Q}$. From Lemma 4.4 we see that $g_N^1(\tau_0) - c_0 = 0$ for some $\tau_0 \in \mathbb{H}$, which yields $Q(g_N^1(\tau_0)) = 0$. Since $P(x)$ is not divisible by $x - c_0$ in $\overline{\mathbb{Q}}(x)$, we have $P(g_N^1(\tau_0)) \neq 0$. It gives a contradiction because $h(\tau)$ is weakly holomorphic. Thus any zero of $Q(x)$ belongs to C_N. Since $Q(x) \in \mathbb{Q}(x)$, we derive that $Q(x)$ has a zero $c \in C_N$ if and only if $f_{N,c}(x) \mid Q(x)$. Therefore, we get $\mathcal{O}_N^1(\mathbb{Q}) \subseteq \mathbb{Q}\big[g_N^1(\tau), f_{N,c}(g_N^1(\tau))^{-1}\big]_{c\in C_N}$. And so, we have the theorem by (4.4).

□

5 Generators of $\mathcal{O}_{1,N}(\mathbb{Q})$

In this section, we shall construct generators of the ring $\mathcal{O}_{1,N}(\mathbb{Q})$ over $\mathbb{Q}$ when N is divisible by $4, 5, 6, 7$ or 9.

Lemma 5.1. *Let m (> 1) be an integer.*

- (i) $f_{\left[\begin{smallmatrix} k/m \\ 0 \end{smallmatrix}\right]}(\tau) \in \mathcal{O}_m^1(\mathbb{Q})$ *for* $k \in \mathbb{Z} \setminus m\mathbb{Z}$.
- (ii) $\mathcal{F}_m^1(\mathbb{Q}) = \mathcal{F}_1(f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau))$.

Proof. By Proposition 3.1 we are certain that $f_{\left[\begin{smallmatrix} k/m \\ 0 \end{smallmatrix}\right]}(\tau)$ is weakly holomorphic and invariant under the action of G_m. Hence it has rational Fourier coefficients by Proposition 2.1. Furthermore, for $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma^1(m)$ we deduce by Proposition 2.1 and 3.1 that

$$\Big(f_{\left[\begin{smallmatrix} k/m \\ 0 \end{smallmatrix}\right]} \circ \gamma\Big)(\tau) = f_{{}^t\gamma\left[\begin{smallmatrix} k/m \\ 0 \end{smallmatrix}\right]}(\tau) = f_{\left[\begin{smallmatrix} ka/m \\ kb/m \end{smallmatrix}\right]}(\tau) = f_{\left[\begin{smallmatrix} k/m \\ 0 \end{smallmatrix}\right]}(\tau)$$

since $a \equiv 1 \pmod{m}$ and $b \equiv 0 \pmod{m}$. Thus $f_{\left[\begin{smallmatrix} k/m \\ 0 \end{smallmatrix}\right]}(\tau)$ is modular for $\Gamma^1(m)$, which proves (i).

By (i) we have $\mathcal{F}_1(f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)) \subseteq \mathcal{F}_m^1(\mathbb{Q})$. Let $\sigma \in \mathrm{Gal}(\mathcal{F}_m/\mathcal{F}_1(f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)))$. Since $f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)$ is G_m-invariant, σ is represented by $\alpha_\sigma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}/m\mathbb{Z})/\{\pm I_2\}$. Then we get

$$f_{\left[\begin{smallmatrix}1/m\\0\end{smallmatrix}\right]}(\tau) = f_{\left[\begin{smallmatrix}1/m\\0\end{smallmatrix}\right]}(\tau)^{\sigma} = f_{{}^t\alpha_\sigma\left[\begin{smallmatrix}1/m\\0\end{smallmatrix}\right]}(\tau) = f_{\left[\begin{smallmatrix}a/m\\b/m\end{smallmatrix}\right]}(\tau). \tag{5.1}$$

Therefore, we achieve $a \equiv d \equiv \pm 1 \pmod{m}$ and $b \equiv 0 \pmod{m}$ by Proposition 3.3 (i), which yields $\mathcal{F}_1(f_{\left[\begin{smallmatrix}1/m\\0\end{smallmatrix}\right]}(\tau)) \supseteq \mathcal{F}_m^1(\mathbb{Q})$ by (4.6). This completes the proof.

□

For integers $m > 3$ and $N > m$ such that $N \equiv 0 \pmod{m}$, let

$$f_{m,N}^1(\tau) = \frac{f_{\left[\begin{smallmatrix}1/N\\0\end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix}1/m\\0\end{smallmatrix}\right]}(\tau)}{f_{\left[\begin{smallmatrix}2/m\\0\end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix}1/m\\0\end{smallmatrix}\right]}(\tau)}.$$

Note that $f_{\left[\begin{smallmatrix}2/m\\0\end{smallmatrix}\right]}(\tau) \neq f_{\left[\begin{smallmatrix}1/m\\0\end{smallmatrix}\right]}(\tau)$ by Proposition 3.3 (i). It then follows from Remark 3.4 and Lemma 5.1 that $f_{m,N}^1(\tau)$ is a modular unit in $\mathcal{O}_N^1(\mathbb{Q})$.

THEOREM 5.2. *Let $m > 3$ and N $(> m)$ be integers such that $N \equiv 0 \pmod{m}$. Then we have*

$$\begin{aligned} \mathcal{O}_{1,N}(\mathbb{Q}) &= \mathcal{O}_{1,m}(\mathbb{Q})\big[f_{m,N}^1(N\tau)\big] \\ \mathcal{O}_N^1(\mathbb{Q}) &= \mathcal{O}_m^1(\mathbb{Q})\big[f_{m,N}^1(\tau)\big]. \end{aligned}$$

PROOF. It suffices to prove the last assertion by the isomorphism (4.4). Clearly, $\mathcal{O}_N^1(\mathbb{Q}) \supseteq \mathcal{O}_m^1(\mathbb{Q})\big[f_{m,N}^1(\tau)\big]$.

As for the converse inclusion, let $h(\tau) \in \mathcal{O}_N^1(\mathbb{Q})$. We see from Lemma 5.1 that

$$\mathcal{F}_N^1(\mathbb{Q}) = \mathcal{F}_1(f_{\left[\begin{smallmatrix}1/N\\0\end{smallmatrix}\right]}(\tau)) = \mathcal{F}_m^1(\mathbb{Q})(f_{m,N}^1(\tau)).$$

Thus $h = h(\tau)$ can be written as

$$h = \sum_{i=0}^{d-1} c_i f^i \tag{5.2}$$

where $f = f_{m,N}^1(\tau)$, $d = [\mathcal{F}_N^1(\mathbb{Q}) : \mathcal{F}_m^1(\mathbb{Q})]$ and $c_i \in \mathcal{F}_m^1(\mathbb{Q})$ for every i. By multiplying both sides of (5.2) by f^j for each $j = 0, 1, \ldots, d-1$ and taking the trace $\mathrm{Tr} = \mathrm{Tr}_{\mathcal{F}_N^1(\mathbb{Q})/\mathcal{F}_m^1(\mathbb{Q})}$, we obtain a linear system

$$\begin{bmatrix} \mathrm{Tr}(1) & \mathrm{Tr}(f) & \cdots & \mathrm{Tr}(f^{d-1}) \\ \mathrm{Tr}(f) & \mathrm{Tr}(f^2) & \cdots & \mathrm{Tr}(f^d) \\ \vdots & \vdots & & \vdots \\ \mathrm{Tr}(f^{d-1}) & \mathrm{Tr}(f^d) & \cdots & \mathrm{Tr}(f^{2d-2}) \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \\ \vdots \\ c_{d-1} \end{bmatrix} = \begin{bmatrix} \mathrm{Tr}(h) \\ \mathrm{Tr}(fh) \\ \vdots \\ \mathrm{Tr}(f^{d-1}h) \end{bmatrix}.$$

For simplicity, put

$$T = \begin{bmatrix} \mathrm{Tr}(1) & \mathrm{Tr}(f) & \cdots & \mathrm{Tr}(f^{d-1}) \\ \mathrm{Tr}(f) & \mathrm{Tr}(f^2) & \cdots & \mathrm{Tr}(f^d) \\ \vdots & \vdots & & \vdots \\ \mathrm{Tr}(f^{d-1}) & \mathrm{Tr}(f^d) & \cdots & \mathrm{Tr}(f^{2d-2}) \end{bmatrix}.$$

Since $f, h \in \mathcal{O}_N^1(\mathbb{Q})$, we attain $c_i \in \det(T)^{-1}\mathcal{O}_m^1(\mathbb{Q})$ for $i = 0, 1, \ldots, d-1$. Now, let $f_1, f_2, \ldots, f_d$ be the Galois conjugates of f over $\mathcal{F}_m^1(\mathbb{Q})$. Then, by the Vandermonde determinant formula, we get

$$\begin{aligned} \det(T) &= \begin{vmatrix} \sum_{k=1}^d f_k^0 & \sum_{k=1}^d f_k^1 & \cdots & \sum_{k=1}^d f_k^{d-1} \\ \sum_{k=1}^d f_k^1 & \sum_{k=1}^d f_k^2 & \cdots & \sum_{k=1}^d f_k^d \\ \vdots & \vdots & & \vdots \\ \sum_{k=1}^d f_k^{d-1} & \sum_{k=1}^d f_k^d & \cdots & \sum_{k=1}^d f_k^{2d-2} \end{vmatrix} \\ &= \begin{vmatrix} f_1^0 & f_2^0 & \cdots & f_d^0 \\ f_1^1 & f_2^1 & \cdots & f_d^1 \\ \vdots & \vdots & & \vdots \\ f_1^{d-1} & f_2^{d-1} & \cdots & f_d^{d-1} \end{vmatrix} \cdot \begin{vmatrix} f_1^0 & f_1^1 & \cdots & f_1^{d-1} \\ f_2^0 & f_2^1 & \cdots & f_2^{d-1} \\ \vdots & \vdots & & \vdots \\ f_d^0 & f_d^1 & \cdots & f_d^{d-1} \end{vmatrix} \\ &= \prod_{1 \le i < j \le d} (f_i - f_j)^2. \end{aligned}$$

It follows from Proposition 3.3, Lemma 5.1 and (5.1) that each $f_i - f_j$ is of the form

$$\frac{f_{\left[\begin{smallmatrix} a/N \\ b/N \end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)}{f_{\left[\begin{smallmatrix} 2/m \\ 0 \end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)} - \frac{f_{\left[\begin{smallmatrix} c/N \\ d/N \end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)}{f_{\left[\begin{smallmatrix} 2/m \\ 0 \end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)} = \frac{f_{\left[\begin{smallmatrix} a/N \\ b/N \end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix} c/N \\ d/N \end{smallmatrix}\right]}(\tau)}{f_{\left[\begin{smallmatrix} 2/m \\ 0 \end{smallmatrix}\right]}(\tau) - f_{\left[\begin{smallmatrix} 1/m \\ 0 \end{smallmatrix}\right]}(\tau)}$$

for some $a, b, c, d \in \mathbb{Z}$ such that $\begin{bmatrix} a/N \\ b/N \end{bmatrix} \not\equiv \pm \begin{bmatrix} c/N \\ d/N \end{bmatrix} \pmod{\mathbb{Z}^2}$ and $\gcd(a, b, N) = \gcd(c, d, N) = 1$. Hence $\det(T)$ is a modular unit in $\mathcal{O}_m^1(\mathbb{Q})$ by Remark 3.4, and so $c_i \in \mathcal{O}_m^1(\mathbb{Q})$ for every i. Therefore, $h(\tau) \in \mathcal{O}_m^1(\mathbb{Q})\big[f_{m,N}^1(\tau)\big]$ as desired. □

Corollary 5.3. *Let N be a positive integer.*

(i) *If $N > 4$ and $N \equiv 0 \pmod 4$, then we have*

$$\begin{aligned} \mathcal{O}_{1,N}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,4}(\tau), g_{1,4}(\tau)^{-1}, (g_{1,4}(\tau) - 16)^{-1}, f_{4,N}^1(N\tau)\big] \\ \mathcal{O}_N^1(\mathbb{Q}) &= \mathbb{Q}\big[g_4^1(\tau), g_4^1(\tau)^{-1}, (g_4^1(\tau) - 16)^{-1}, f_{4,N}^1(\tau)\big]. \end{aligned}$$

(ii) *If $N > 5$ and $N \equiv 0 \pmod 5$, then we have*

$$\begin{aligned} \mathcal{O}_{1,N}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,5}(\tau), g_{1,5}(\tau)^{-1}, (g_{1,5}(\tau)^2 - 11g_{1,5}(\tau) - 1)^{-1}, f_{5,N}^1(N\tau)\big] \\ \mathcal{O}_N^1(\mathbb{Q}) &= \mathbb{Q}\big[g_5^1(\tau), g_5^1(\tau)^{-1}, (g_5^1(\tau)^2 - 11g_5^1(\tau) - 1)^{-1}, f_{5,N}^1(\tau)\big]. \end{aligned}$$

(iii) *If* $N > 6$ *and* $N \equiv 0 \pmod 6$*, then we have*

$$\begin{aligned}
\mathcal{O}_{1,N}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,6}(\tau), g_{1,6}(\tau)^{-1}, (g_{1,6}(\tau)-8)^{-1}, (g_{1,6}(\tau)+1)^{-1}, f^1_{6,N}(N\tau)\big] \\
\mathcal{O}^1_{N}(\mathbb{Q}) &= \mathbb{Q}\big[g^1_{6}(\tau), g^1_{6}(\tau)^{-1}, (g^1_{6}(\tau)-8)^{-1}, (g^1_{6}(\tau)+1)^{-1}, f^1_{6,N}(\tau)\big].
\end{aligned}$$

(iv) *If* $N > 7$ *and* $N \equiv 0 \pmod 7$*, then we have*

$$\begin{aligned}
\mathcal{O}_{1,N}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,7}(\tau), g_{1,7}(\tau)^{-1}, (g_{1,7}(\tau)-1)^{-1}, (g_{1,7}(\tau)^3-8g_{1,7}(\tau)^2+5g_{1,7}(\tau)+1)^{-1}, \\
&\qquad f^1_{7,N}(N\tau)\big] \\
\mathcal{O}^1_{N}(\mathbb{Q}) &= \mathbb{Q}\big[g^1_{7}(\tau), g^1_{7}(\tau)^{-1}, (g^1_{7}(\tau)-1)^{-1}, (g^1_{7}(\tau)^3-8g^1_{7}(\tau)^2+5g^1_{7}(\tau)+1)^{-1}, f^1_{7,N}(\tau)\big].
\end{aligned}$$

(v) *If* $N > 9$ *and* $N \equiv 0 \pmod 9$*, then we have*

$$\begin{aligned}
\mathcal{O}_{1,N}(\mathbb{Q}) &= \mathbb{Q}\big[g_{1,9}(\tau), g_{1,9}(\tau)^{-1}, (g_{1,9}(\tau)-1)^{-1}, (g_{1,9}(\tau)^2-g_{1,9}(\tau)+1)^{-1}, \\
&\qquad (g_{1,9}(\tau)^3-6g_{1,9}(\tau)^2+3g_{1,9}(\tau)+1)^{-1}, f^1_{9,N}(N\tau)\big] \\
\mathcal{O}^1_{N}(\mathbb{Q}) &= \mathbb{Q}\big[g^1_{9}(\tau), g^1_{9}(\tau)^{-1}, (g^1_{9}(\tau)-1)^{-1}, (g^1_{9}(\tau)^2-g^1_{9}(\tau)+1)^{-1}, \\
&\qquad (g^1_{9}(\tau)^3-6g^1_{9}(\tau)^2+3g^1_{9}(\tau)+1)^{-1}, f^1_{9,N}(\tau)\big].
\end{aligned}$$

PROOF. It is immediate from Theorem 4.5 and 5.2. □

REMARK 5.4. In order to construct $\mathcal{O}_{1,N}(\mathbb{Q})$ for arbitrary N, it suffices to find generators of the ring $\mathcal{O}_{1,p}(\mathbb{Q})$ for an odd prime $p \geq 11$.

6 Fricke families of level N

For an integer N (> 1), let

$$\mathcal{R}_N = \{\mathbf{r} \in \mathbb{Q}^2 \mid \mathbf{r} \text{ has the primitive denominator } N\}.$$

A family $\{h_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N}$ of functions in $\mathcal{F}_N$ is called a *Fricke family* of level N if it satisfies the following conditions:

(i) Each $h_{\mathbf{r}}(\tau)$ is weakly holomorphic.

(ii) $h_{\mathbf{r}}(\tau)$ depends only on $\pm\mathbf{r} \pmod{\mathbb{Z}^2}$.

(iii) $h_{\mathbf{r}}(\tau)^\alpha = h_{{}^t\alpha\mathbf{r}}(\tau)$ for all $\alpha \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$.

Note that the set of Fricke functions $\{f_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N}$ is a Fricke family of level N by Proposition 3.1. In this section, by $\mathrm{Fr}(N)$ we mean the set of all Fricke families of level N. Then one can endow this set the natural ring structure:

$$\begin{aligned}\{h_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N}+\{k_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N} &= \{(h_{\mathbf{r}}+k_{\mathbf{r}})(\tau)\}_{\mathbf{r}\in\mathcal{R}_N}\\ \{h_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N}\cdot\{k_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N} &= \{(h_{\mathbf{r}}k_{\mathbf{r}})(\tau)\}_{\mathbf{r}\in\mathcal{R}_N}.\end{aligned}$$

Eum et al ([2]) classify all Fricke families of level N when $N\equiv 0 \pmod 4$ by using the relation between $\mathrm{Fr}(N)$ and $\mathcal{O}^1_N(\mathbb{Q})$ as follows:

Proposition 6.1. *For an integer $N>1$, the map*

$$\begin{aligned}\mathrm{Fr}(N) &\longrightarrow \mathcal{O}^1_N(\mathbb{Q})\\ \{h_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N} &\longmapsto h_{\left[\begin{smallmatrix}1/N\\0\end{smallmatrix}\right]}(\tau)\end{aligned}$$

is a well-defined ring isomorphism.

Proof. [2, Theorem 4.3]. □

Theorem 6.2. *Let $N>1$ be a positive integer. Then a family $\{h_{\mathbf{r}}(\tau)\}_{\mathbf{r}\in\mathcal{R}_N}$ of functions in $\mathcal{F}_N$ is a Fricke family of level N if and only if the following condition holds:*

(i) *When $N\equiv 0 \pmod 4$, there exists a polynomial $P(x_1,x_2,x_3,x_4)\in\mathbb{Q}[x_1,x_2,x_3,x_4]$ such that*

$$h_{\mathbf{r}}(\tau)=P\big(\mathfrak{g}_{\mathbf{r}}(\tau),\mathfrak{g}_{\mathbf{r}}(\tau)^{-1},(\mathfrak{g}_{\mathbf{r}}(\tau)-16)^{-1},\mathfrak{f}_{\mathbf{r}}(\tau)\big)$$

for all $\mathbf{r}\in\mathcal{R}_N$, where

$$\mathfrak{g}_{\mathbf{r}}(\tau)=\frac{g_{(N/2)\mathbf{r}}(\tau)^8}{g_{(N/4)\mathbf{r}}(\tau)^8}\quad \text{and}\quad \mathfrak{f}_{\mathbf{r}}(\tau)=\frac{f_{\mathbf{r}}(\tau)-f_{(N/4)\mathbf{r}}(\tau)}{f_{(N/2)\mathbf{r}}(\tau)-f_{(N/4)\mathbf{r}}(\tau)}.$$

(ii) *When $N\equiv 0 \pmod 5$, there exists a polynomial $P(x_1,x_2,x_3,x_4)\in\mathbb{Q}[x_1,x_2,x_3,x_4]$ such that*

$$h_{\mathbf{r}}(\tau)=P\big(\mathfrak{g}_{\mathbf{r}}(\tau),\mathfrak{g}_{\mathbf{r}}(\tau)^{-1},(\mathfrak{g}_{\mathbf{r}}(\tau)^2-11\mathfrak{g}_{\mathbf{r}}(\tau)-1)^{-1},\mathfrak{f}_{\mathbf{r}}(\tau)\big)$$

for all $\mathbf{r}\in\mathcal{R}_N$, where

$$\mathfrak{g}_{\mathbf{r}}(\tau)=\frac{g_{(2N/5)\mathbf{r}}(\tau)^5}{g_{(N/5)\mathbf{r}}(\tau)^5}\quad \text{and}\quad \mathfrak{f}_{\mathbf{r}}(\tau)=\frac{f_{\mathbf{r}}(\tau)-f_{(N/5)\mathbf{r}}(\tau)}{f_{(2N/5)\mathbf{r}}(\tau)-f_{(N/5)\mathbf{r}}(\tau)}.$$

(iii) *When $N\equiv 0 \pmod 6$, there exists a polynomial $P(x_1,x_2,x_3,x_4,x_5)\in\mathbb{Q}[x_1,x_2,x_3,x_4,x_5]$ such that*

$$h_{\mathbf{r}}(\tau)=P\big(\mathfrak{g}_{\mathbf{r}}(\tau),\mathfrak{g}_{\mathbf{r}}(\tau)^{-1},(\mathfrak{g}_{\mathbf{r}}(\tau)-8)^{-1},(\mathfrak{g}_{\mathbf{r}}(\tau)+1)^{-1},\mathfrak{f}_{\mathbf{r}}(\tau)\big)$$

for all $\mathbf{r}\in\mathcal{R}_N$, where

$$\mathfrak{g}_{\mathbf{r}}(\tau)=\frac{g_{(N/2)\mathbf{r}}(\tau)^3}{g_{(N/6)\mathbf{r}}(\tau)^3}\quad \text{and}\quad \mathfrak{f}_{\mathbf{r}}(\tau)=\frac{f_{\mathbf{r}}(\tau)-f_{(N/6)\mathbf{r}}(\tau)}{f_{(N/3)\mathbf{r}}(\tau)-f_{(N/6)\mathbf{r}}(\tau)}.$$

(iv) *When* $N \equiv 0 \pmod 7$, *there exists a polynomial* $P(x_1,x_2,x_3,x_4,x_5) \in \mathbb{Q}[x_1,x_2,x_3,x_4,x_5]$ *such that*

$$h_{\mathbf{r}}(\tau) = P\big(\mathfrak{g}_{\mathbf{r}}(\tau), \mathfrak{g}_{\mathbf{r}}(\tau)^{-1}, (\mathfrak{g}_{\mathbf{r}}(\tau)-1)^{-1}, (\mathfrak{g}_{\mathbf{r}}(\tau)^3 - 8\mathfrak{g}_{\mathbf{r}}(\tau)^2 + 5\mathfrak{g}_{\mathbf{r}}(\tau)+1)^{-1}, \mathfrak{f}_{\mathbf{r}}(\tau)\big)$$

for all $\mathbf{r} \in \mathcal{R}_N$, *where*

$$\mathfrak{g}_{\mathbf{r}}(\tau) = \frac{g_{(2N/7)\mathbf{r}}(\tau)^2 g_{(3N/7)\mathbf{r}}(\tau)}{g_{(N/7)\mathbf{r}}(\tau)^3} \quad \text{and} \quad \mathfrak{f}_{\mathbf{r}}(\tau) = \frac{f_{\mathbf{r}}(\tau) - f_{(N/7)\mathbf{r}}(\tau)}{f_{(2N/7)\mathbf{r}}(\tau) - f_{(N/7)\mathbf{r}}(\tau)}.$$

(v) *When* $N \equiv 0 \pmod 9$, *there exists a polynomial* $P(x_1,x_2,x_3,x_4,x_5,x_6) \in \mathbb{Q}[x_1,x_2,x_3,x_4,x_5,x_6]$ *such that*

$$\begin{aligned} h_{\mathbf{r}}(\tau) &= P\big(\mathfrak{g}_{\mathbf{r}}(\tau), \mathfrak{g}_{\mathbf{r}}(\tau)^{-1}, (\mathfrak{g}_{\mathbf{r}}(\tau)-1)^{-1}, (\mathfrak{g}_{\mathbf{r}}(\tau)^2 - \mathfrak{g}_{\mathbf{r}}(\tau)+1)^{-1}, \\ &\qquad (\mathfrak{g}_{\mathbf{r}}(\tau)^3 - 6\mathfrak{g}_{\mathbf{r}}(\tau)^2 + 3\mathfrak{g}_{\mathbf{r}}(\tau)+1)^{-1}, \mathfrak{f}_{\mathbf{r}}(\tau)\big) \end{aligned}$$

for all $\mathbf{r} \in \mathcal{R}_N$, *where*

$$\mathfrak{g}_{\mathbf{r}}(\tau) = \frac{g_{(2N/9)\mathbf{r}}(\tau) g_{(4N/9)\mathbf{r}}(\tau)}{g_{(N/9)\mathbf{r}}(\tau)^2} \quad \text{and} \quad \mathfrak{f}_{\mathbf{r}}(\tau) = \frac{f_{\mathbf{r}}(\tau) - f_{(N/9)\mathbf{r}}(\tau)}{f_{(2N/9)\mathbf{r}}(\tau) - f_{(N/9)\mathbf{r}}(\tau)}.$$

PROOF. For $\mathbf{r} = \begin{bmatrix} a/N \\ b/N \end{bmatrix} \in \mathcal{R}_N$, we can take $\alpha_{\mathbf{r}} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ with $c, d \in \mathbb{Z}$. For each $m = 4, 5, 6, 7$ and 9, if $N > m$ and $N \equiv 0 \pmod m$, then we achieve by Proposition 3.2 (i) that

$$\begin{aligned} g_m^1(\tau)^{\alpha_{\mathbf{r}}} &= \mathfrak{g}_{\mathbf{r}}(\tau) \\ f_{m,N}^1(\tau)^{\alpha_{\mathbf{r}}} &= \mathfrak{f}_{\mathbf{r}}(\tau). \end{aligned}$$

Here we note that $\mathfrak{f}_{\mathbf{r}}(\tau) = 0$ if $N = m$. Then the theorem is an immediate consequence of Theorem 4.5, Corollary 5.3 and Proposition 6.1.

□

References

[1] D. A. Cox, *Primes of the Form $x^2 + ny^2$: Fermat, Class Field, and Complex Multiplication*, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1989.

[2] I. S. Eum, J. K. Koo and D. H. Shin, *Determination of the Fricke families*, arXiv:1501.04193, 2015.

[3] K. Harada, *"Moonshine" of Finite Groups*, EMS Series of Lectures in Mathematics, European Mathematical Society (EMS), Zürich, 2010.

[4] J. K. Koo and D. H. Shin, *On some arithmetic properties of Siegel functions*, Math. Z. 264 (2010), 137–177.

[5] D. Kubert and S. Lang, *Modular Units*, Grundlehren der mathematischen Wissenschaften 244, Spinger-Verlag, New York-Berlin, 1981.

[6] S. Lang, *Elliptic Functions*, 2nd ed., Spinger-Verlag, New York, 1987.

[7] C. L. Siegel, *Lectures on advanced analytic number theory*, Tata Institute of Fundamental Research Lectures on Mathematics, 23, Tata Institute of Fundamental Research, Bombay, 1965.

[8] Y. Yang, *Transformation formulas for generalized Dedekind eta functions*, Bull. London Math. Soc. 36 (2004), 671–682.

Ja Kyung Koo
Department of Mathematical Sciences
KAIST
Daejeon 305-701
Republic of Korea

E-mail address: jkkoo@math.kaist.ac.kr

Dong Sung Yoon
National Institute for Mathematical Sciences
Daejeon 305-811
Republic of Korea

E-mail address: dsyoon@nims.re.kr